

RANDOM WALKS AND THE POROUS MEDIUM EQUATION

C. CORTÁZAR, M. ELGUETA, S. MARTÍNEZ AND J. D. ROSSI

ABSTRACT. In this paper we consider a fixed initial condition u_0 and we study the limit as $\varepsilon \rightarrow 0$ of u_ε , the solution to the re-scaled problem

$$u_t(x, y) = \frac{1}{\varepsilon^2} \left(\int_R J \left(\frac{x-y}{\varepsilon u(y, t)} \right) \frac{dy}{\varepsilon} - u(x, t) \right) \quad \text{in } R \times [0, \infty)$$

with initial condition $u_\varepsilon(x, 0) = u_0(x)$. Here J is a smooth non negative even function supported in the interval $[-1, 1]$. Moreover it is assumed that $\int J = 1$ and that J is decreasing on $[0, 1]$.

We prove that, under adequate hypothesis on the initial condition, the limit

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon = u$$

is the solution to the well known porous medium equation $v_t = D(v^3)_{xx}$, with initial condition $u_\varepsilon(x, 0) = u_0(x)$ for a suitable constant D .

INTRODUCTION.

Let $J : \mathbb{R}^N \rightarrow \mathbb{R}$ be a nonnegative, smooth function such that $\int_{\mathbb{R}^N} J(y) dy = 1$. Equations of the form

$$u_t(x, t) = \int_{\mathbb{R}^N} J(x-y)u(y, t) dy - u(x, t) \quad \text{in } \mathbb{R}^N \times [0, \infty) \quad (1.1)$$

have been widely used to model diffusion processes, see [2], [4], [10], [11], [12]. As stated in [10] if $u(x, t)$ is thought of as a density at the point x at time t and $J(x-y)$ is thought of as the probability distribution of jumping from location y to location x , then $\int_{\mathbb{R}^N} J(x-y)u(y, t) dy$ is the rate at which individuals are arriving to position x from all other places. On the other hand $-u(x, t) = -\int_{\mathbb{R}^N} J(x-y)u(x, t) dy$ is the rate at which they are leaving location x to travel to all other sites. This consideration, in the absence of external sources, leads immediately to the fact that the density u satisfies equation (1.1).

Throughout this note we shall assume also that J is a decreasing radial function whose support is the unit ball. Under these hypotheses we have that individuals that are at a position x are not allowed, with probability 1, to jump to a locations

Key words and phrases. Nonlocal diffusion, free boundaries.

C. Cortázar and M. Elgueta partially supported by FONDECYT 1070944. S. Martínez partially supported by FONDECYT 1090183 and FONDAP de Matemáticas Aplicada. J. D. Rossi partially supported by MTM2004-02223, MEC, Spain, by UBA X066 and by CONICET, Argentina.

y such that $\|x - y\| > 1$. In this fashion we like to think of this process as a, continuous in time, random walk of step's size 1.

Another equation that has been widely used to model diffusion processes is the classical heat equation

$$v_t = D\Delta v. \tag{1.2}$$

Equations (1.1) and (1.2) share several properties. For example the initial value problem is well posed for suitable initial data and a maximum principle, and hence a comparison one, holds for both of them.

An important relation between equations (1.1) and (1.2) is that solutions of a properly rescaled version of (1.2) converge to the solution of (1.1) when the scaling parameter tends to zero. More precisely, let $\varepsilon > 0$ and consider the rescaled equation

$$\varepsilon^2 u_t(x, t) = \int_{\mathbb{R}^N} J\left(\frac{x - y}{\varepsilon}\right) \frac{u(y, t)}{\varepsilon^N} dy - u(x, t). \tag{1.3}$$

We observe that solutions to (1.3) are random walks, in the sense explained above, where the probability density of jumping from position y to position x is given by

$$J\left(\frac{x - y}{\varepsilon}\right) \frac{1}{\varepsilon^N}$$

so in this case the size of the step of the random walk is ε . On the other hand the ε^2 that appears in the left hand side of (1.3) stands for an increment in the number of steps per unit of time.

The following result is classic:

Theorem 1.1. *Let u_ε be a solution of (1.3) and let v be a solution of (1.2). Assume that*

$$u_\varepsilon(x, 0) = v(x, 0) \in L^1(\mathbb{R}^N),$$

then

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon(x, t) = v(x, t)$$

in $L^1(\mathbb{R}^N \times [0, T])$ for any $T > 0$.

The aim of this note is to exhibit a similar relation between solutions of the porous medium equation, $v_t = \Delta(v^m)$, and suitable random walks. Since solutions v of the porous medium equation have the property that if $v(\cdot, 0)$ is compactly supported in \mathbb{R}^N then so is $v(\cdot, t)$ for all $t > 0$, see [1], we would like to approximate them by random walks that also have this property. Such a random walk was constructed in [6].

OUR RESULT.

The well known porous medium equation

$$v_t = D\Delta(v^m) \text{ in } \mathbb{R}^N \times [0, \infty), \tag{1.4}$$

where $m > 1$ and $D > 0$ is any constant, has been also widely used to model diffusion.

Solutions to (1.4) and (1.2) share several properties such as the fact that the initial value problem is well posed for suitable initial data and a comparison principle holds for both of them. However there exists quite a difference in their behavior as we explain now. While solutions to (1.2) do have the so called infinite speed of propagation of disturbances, this means that if u is a non trivial solution of (1.2) with $u(x, 0) \geq 0$ then $u(x, t) > 0$ for all (x, t) with $t > 0$, solutions of (1.4) do not. See [1]. It is actually proved that if $u \geq 0$ is a solution of (1.4) such that the support of $u(\cdot, 0)$ is contained in a finite interval $[-a, a]$, then there exists a non decreasing function $s : [0, \infty) \rightarrow [0, \infty)$ such that $u(\cdot, t)$ is supported inside $[-s(t), s(t)]$.

At this point it is worth to note here that solutions of (1.3) do have infinite speed of propagation of disturbances. This is no difficult to prove and is left as an exercise.

In [6] a random walk was proposed, in the case $N = 1$, which has the property that if the initial condition has compact support, then the spacial support of the solution is compact for any $t > 0$. This was achieved by considering functions of the form

$$J\left(\frac{x-y}{u(y,t)}\right) \frac{1}{u(y,t)}$$

to represent the probability of jumping from position y to position x .

In this case the random walk, continuous in time, is regulated by the equation

$$u_t(x, y) = \int_{\mathbb{R}} J\left(\frac{x-y}{u(y,t)}\right) dy - u(x, t) \quad \text{in } \mathbb{R} \times [0, \infty). \quad (1.5)$$

Existence, uniqueness and a comparison principle for (1.5) have been proved in [6]. See [3] for the case of arbitrary dimension, that is, in \mathbb{R}^N .

We are now in a position to describe the main result of this note. To do this let an initial condition u_0 be given and let $\varepsilon > 0$. Consider the re-scaled problem

$$u_t(x, y) = \frac{1}{\varepsilon^2} \left(\int_{\mathbb{R}} J\left(\frac{x-y}{\varepsilon u(y,t)}\right) \frac{dy}{\varepsilon} - u(x, t) \right) \quad \text{in } \mathbb{R} \times [0, \infty). \quad (1.6)$$

and let u_ε be a solution of (1.6) with initial condition $u_\varepsilon(x, 0) = u_0(x)$.

We want to determine the escaled limit

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon.$$

Unfortunately at this point we have to make an extra assumption that we are sure it can be removed. The extra assumption is

$$u_\varepsilon(x, 0) = w_0(x) + \delta$$

where $w_0(x)$ is a non negative compactly supported C^∞ function and $\delta > 0$.

We need first the following definition. For $\varepsilon > 0$ and a positive bounded function h we define

$$T_\varepsilon h(x) = \int_{\mathbb{R}} J\left(\frac{y-x}{\varepsilon h(y)}\right) \frac{dy}{\varepsilon} - h(x).$$

We also need the following lemma.

Lemma 1.1. *Let $\delta > 0$ and let h be a smooth function such that $h(x) \geq \delta$. Then*

$$T_\varepsilon h(x) = C(h^3)_{xx}\varepsilon^2 + R(x, \varepsilon)$$

where $R(\cdot, \varepsilon) \in L^1(\mathbb{R})$ and $C = \int J(z)z^2 dz$. Moreover

$$\|R(\cdot, \varepsilon)\|_1 \leq K \frac{\varepsilon^3}{\delta^6}$$

where K depends on $J, \|h\|_\infty, \|h'\|_\infty, \|h''\|_\infty, \|h'''\|_\infty, \|h'\|_1, \|h''\|_1$ and $\|h'''\|_1$ but is independent of ε, δ and $\|h\|_1$.

Proof: The proof consists of expanding the function $T_\varepsilon h(x)$ in a Taylor series in ε about the point $\varepsilon = 0$.

Performing the change of variables $z = \frac{y-x}{\varepsilon}$ we get

$$T_\varepsilon h(x) = \int_{\mathbb{R}} J\left(\frac{z}{h(x+\varepsilon z)}\right) dz - h(x).$$

Hence

$$T_0 h(x) = 0. \tag{1.7}$$

Now

$$\frac{d}{d\varepsilon} T_\varepsilon h(x) = - \int_{\mathbb{R}} J'\left(\frac{z}{h(x+\varepsilon z)}\right) z^2 \frac{h'(x+\varepsilon z)}{h^2(x+\varepsilon z)} dz.$$

and using the fact that J' is an odd function we get

$$\frac{d}{d\varepsilon} T_\varepsilon h(x)|_{\varepsilon=0} = 0. \tag{1.8}$$

Differentiating with respect to ε one more time we obtain

$$\begin{aligned} \frac{d^2}{d\varepsilon^2} T_\varepsilon h(x) &= \int_{\mathbb{R}} J''\left(\frac{z}{h(x+\varepsilon z)}\right) z^4 \left(\frac{h'(x+\varepsilon z)}{h^2(x+\varepsilon z)}\right)^2 dz \\ &\quad - \int_{\mathbb{R}} J'\left(\frac{z}{h(x+\varepsilon z)}\right) z^3 \left(\frac{h'(x+\varepsilon z)}{h^2(x+\varepsilon z)}\right)' dz. \end{aligned}$$

Thus

$$\begin{aligned} \frac{d^2}{d\varepsilon^2} T_\varepsilon h(x)|_{\varepsilon=0} &= \int_{\mathbb{R}} J''\left(\frac{z}{h(x)}\right) z^4 dz \left(\frac{h'(x)}{h^2(x)}\right)^2 \\ &\quad - \int_{\mathbb{R}} J'\left(\frac{z}{h(x)}\right) z^3 dz \left(\frac{h'(x)}{h^2(x)}\right)' \\ &= \int_{\mathbb{R}} J''(w)w^4 dw h(x)(h'(x))^2 - \int_{\mathbb{R}} J'(w)w^3 dw h^4(x) \left(\frac{h'(x)}{h^2(x)}\right)'. \end{aligned}$$

But

$$\int_{\mathbb{R}} J''\left(\frac{z}{h(x)}\right) z^4 dz = -4 \int_{\mathbb{R}} J'\left(\frac{z}{h(x)}\right) z^3 dz = 12 \int_{\mathbb{R}} J\left(\frac{z}{h(x)}\right) z^2 dz$$

and

$$\int_{\mathbb{R}} J'\left(\frac{z}{h(x)}\right) z^3 dz = -3 \int_{\mathbb{R}} J\left(\frac{z}{h(x)}\right) z^2 dz.$$

So setting

$$C = \int_{\mathbb{R}} J(z)z^2 dz$$

we get

$$\begin{aligned} \frac{d^2}{d\varepsilon^2} T_\varepsilon h(x)|_{\varepsilon=0} &= C \left(12h(x)(h'(x))^2 + 3h^4(x) \left(\frac{h'(x)}{h^2(x)} \right)' \right) \\ &= C(6h(x)(h'(x))^2 + 3h^2(x)h''(x)) = C(h^3(x))''. \end{aligned}$$

Or

$$\frac{d^2}{d\varepsilon^2} T_\varepsilon h(x)|_{\varepsilon=0} = C(h^3(x))''. \tag{1.9}$$

Using (1.7), (1.8) and (1.9) we obtain, by Taylor’s theorem,

$$T_\varepsilon h(x) = C(h^3)_{xx}\varepsilon^2 + R(x, \varepsilon)$$

where the remainder can be written as

$$R(x, \varepsilon) = \frac{1}{2} \int_0^\varepsilon \frac{d^3}{ds^3} T_s h(x)(\varepsilon - s)^2 ds.$$

Differentiating the formula for $\frac{d^2}{d\varepsilon^2} T_\varepsilon h(x)$ one more time with respect to ε and plugging in $\frac{d^3}{ds^3} T_s h(x)$ into the above integral the desired estimate is obtained after a straightforward calculation that uses the fact that J is compactly supported. \square

We will give now a proof of our result. We recall that we are assuming that the initial condition is of the form $u_0(x) = w_0(x) + \delta$ with $\delta > 0$ and w_0 a compactly supported non negative C^∞ function.

Theorem 1.2. *Let $w_0 \in C^\infty(\mathbb{R})$ be a nonnegative compactly supported function and let $\delta > 0$. For each $\varepsilon > 0$ let u_ε be the solution (1.6) with initial condition $u_\varepsilon(x, 0) = u_0(x) = w_0(x) + \delta$. Let v be a solution of*

$$v_t = C(v^3)_{xx}$$

with initial condition $v(x, 0) = u_0(x) = w_0(x) + \delta$.

Then

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon = v$$

in $L^1(\mathbb{R} \times [0, \infty])$ for each $T > 0$.

Proof: Existence and uniqueness of the solutions u_ε can be found in [6]. By Lemma 1.1 we can write

$$v_t(x, t) = \frac{1}{\varepsilon^2} \left(\int_{\mathbb{R}} J \left(\frac{y - x}{\varepsilon v(y, t)} \right) \frac{dy}{\varepsilon} - v(x, t) \right) + \frac{1}{\varepsilon^2} R(x, t, \varepsilon)$$

where, due to our hypotheses on w_0 ,

$$\|R(\cdot, t, \varepsilon)\|_1 \leq K \frac{\varepsilon^3}{\delta^6} \text{ for all } t \in [0, T]$$

with K independent of ε and δ .

Let now \underline{w} y \overline{w} solutions of

$$\underline{w}_t(x, t) = \frac{1}{\varepsilon^2} \left(\int_{\mathbb{R}} J \left(\frac{y-x}{\varepsilon \underline{w}(y, t)} \right) \frac{dy}{\varepsilon} - \underline{w}(x, t) \right) - \frac{1}{\varepsilon^2} |R(x, t, \varepsilon)|$$

and

$$\overline{w}_t(x, t) = \frac{1}{\varepsilon^2} \left(\int_{\mathbb{R}} J \left(\frac{y-x}{\varepsilon \overline{w}(y, t)} \right) \frac{dy}{\varepsilon} - \overline{w}(x, t) \right) + \frac{1}{\varepsilon^2} |R(x, t, \varepsilon)|$$

with initial data

$$\underline{w}(x, 0) = \overline{w}(x, 0) = u_0(x) + \delta,$$

respectively.

Clearly, by comparison,

$$\underline{w}(x, t) \leq v(x, t) \leq \overline{w}(x, t),$$

and

$$\underline{w}(x, t) \leq u_\varepsilon(x, t) \leq \overline{w}(x, t).$$

Now since the quantity inside the double integral is non negative, in this step we are using strongly the hypothesis that J is decreasing in $[0, 1]$, we can apply Fubini's theorem and get

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} (\overline{w} - \underline{w})(x, t) dx &= \\ &= \int_{\mathbb{R}} \frac{1}{\varepsilon^2} \left(\int_{\mathbb{R}} J \left(\frac{y-x}{\varepsilon \overline{w}(y, t)} \right) \frac{dy}{\varepsilon} - \overline{w}(x, t) \right) dx \\ &\quad - \int_{\mathbb{R}} \frac{1}{\varepsilon^2} \left(\int_{\mathbb{R}} J \left(\frac{y-x}{\varepsilon \underline{w}(y, t)} \right) \frac{dy}{\varepsilon} - \underline{w}(x, t) \right) dx \\ &\quad + 2 \frac{1}{\varepsilon^2} \int_{\mathbb{R}} |R(x, t, \varepsilon)| dx \\ &= 2 \frac{1}{\varepsilon^2} \int_{\mathbb{R}} |R(x, t, \varepsilon)| dx \leq 2K \frac{\varepsilon}{\delta^6}. \end{aligned}$$

This implies

$$\|\overline{w} - \underline{w}\|_1 \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

and hence

$$u_\varepsilon \rightarrow v$$

en $L^1(\mathbb{R} \times [0, T])$ for any $T > 0$ as $\varepsilon \rightarrow 0$ as we wanted to prove. \square

We end this note with the following remark: the extra hypothesis $u_\varepsilon(x, 0) = v(x, 0) = w_0(x) + \delta$ with $\delta > 0$ in Theorem 1.2 has to be replaced by $u_\varepsilon(x, 0) = v(x, 0) = w_0(x)$ with $w_0 \in L^1(\mathbb{R})$. We hope we can do this in the near future.

Once this is done an interesting question arises: Assume that w_0 is even with respect to the origin and $\text{supp } w_0 = [-a, a]$. Then it is known that there exist non decreasing functions, called the free boundaries, $s : [0, \infty) \rightarrow [0, \infty)$ and $s_\varepsilon : [0, \infty) \rightarrow [0, \infty)$ such that $\text{supp } v(\cdot, t) = [-s(t), s(t)]$ and $\text{supp } u_\varepsilon(\cdot, t) = [-s_\varepsilon(t), s_\varepsilon(t)]$. A natural question to ask is whether and in which sense $\lim_{\varepsilon \rightarrow 0} s_\varepsilon = s$.

REFERENCES

- [1] D. G. Aronson. *The porous medium equation* A. Fasano and M. Primicerio eds. Lecture Notes in Math. 1224. Springer Verlag. (1986).
- [2] P. Bates, P. Fife, X. Ren and X. Wang. *Traveling waves in a convolution model for phase transitions*. Arch. Rat. Mech. Anal., 138, 105-136, (1997).
- [3] M. Bogoya. *A nonlocal nonlinear diffusion equation in higher space dimensions*. J. Math. Anal. Appl. 344 (2008), no. 2, 601–615. (2008).
- [4] X. Chen. *Existence, uniqueness and asymptotic stability of travelling waves in nonlocal evolution equations*. Adv. Differential Equations, 2, 125-160, (1997).
- [5] C. Cortázar, J. Coville, M. Elgueta and S. Martínez. *A non local inhomogeneous dispersal process*. J. Differential Equations 241 (2007), no. 2, 332-358.
- [6] C. Cortázar, M. Elgueta and J. Rossi. *A non-local diffusion equation whose solutions develop a free boundary*. Ann. Inst. H. Poincaré 6 (2) (2005) 269-281.
- [7] C. Cortázar, M. Elgueta and J. Rossi. *Nonlocal diffusion problems that approximate the heat equation with Dirichlet boundary conditions*. Israel J. Math. 170 (2009), 53–60.
- [8] C. Cortázar, M. Elgueta, J. Rossi and N. Wolanski. *Boundary fluxes for non-local diffusion*. J. Differential Equations 234 (2007) 360-390.
- [9] C. Cortázar, M. Elgueta, J. Rossi and N. Wolanski. *How to approximate the heat equation with Neumann boundary conditions by nonlocal diffusion problems*. Archive for Rational Mechanics and Analysis. Vol. 187 (1),(2008) 137-156.
- [10] P. Fife. *Some nonclassical trends in parabolic and parabolic-like evolutions*. Trends in nonlinear analysis, 153–191, Springer, Berlin, 2003.
- [11] C. Lederman and N. Wolanski. *A free boundary problem from nonlocal combustion*. Preprint
- [12] X. Wang. *Meta-stability and stability of patterns in a convolution model for phase transitions*. Preprint.

C. Cortázar

Facultad de Matemáticas
P. Universidad Católica de Chile
Vicuña Mackenna 4860, Santiago, Chile

M. Elgueta

Facultad de Matemáticas
P. Universidad Católica de Chile
Vicuña Mackenna 4860, Santiago, Chile

Salomé Martínez

Departamento de Ingeniería Matemática
and Centro de Modelamiento Matemático, UMI 2807 CNRS-UChile
Universidad de Chile, Casilla 170 Correo 3, Santiago, Chile.

Julio D. Rossi

Depto. Matemática, FCEyN,
Universidad de Buenos Aires
Ciudad Universitaria, Pab 1,
(1428), Buenos Aires, Argentina.

Recibido: 1 de agosto de 2008

Aceptado: 2 de marzo de 2009